arXiv:1207.4256v1 [quant-ph] 18 Jul 2012

Dynamics and thermodynamics of linear quantum open systems

Esteban Martínez[1] and Juan Pablo Paz[1, 2]
[1]*Departamento de Física, FCEyN, UBA, Ciudad Universitaria Pabellón 1, 1428 Buenos Aires, Argentina*
[2]*IFIBA CONICET, UBA, FCEyN, UBA, Ciudad Universitaria Pabellón 1, 1428 Buenos Aires, Argentina*

(Dated: July 19, 2012)

We analyze the behavior of a network of quantum oscillators coupled with a number of external environments. We show that the dynamics is such that the quantum state of the network always obeys a local master equation with a simple analytical solution. We use this to study the emergence of thermodynamical laws in the stationary regime, achieved for sufficiently long times if the environments are dissipative. We show that the validity of the second law implies the impossibility of building a quantum refrigerator without moving parts (therefore, a quantum absorption refrigerators requires non-linearity as an crucial ingredient, as recently proposed by Kosloff and others citeKosloff1,Kosloff2). We also show that the third law imposes strong constraints on the low frequency behavior of the environmental spectral densities.

PACS numbers: 03.65.Yz, 03.67.Bg, 37.10.Rs

Understanding the origin of classical physics from quantum laws is important not only for fundamental but also for practical reasons [1]. In fact, more and more experimentally accesible systems manifest quantum effects while having a large number of degrees of freedom. In such systems, many of which are being developed as part of the quest for the quantum computer [2], it is of fundamental importance to study the emergence of thermodynamical laws [3–5] to understand the dynamics of heat transport and decoherence. We present here some general results concerning not only the evolution of linear open quantum networks but also the emergence of thermodynamics in such systems. This may be relevant not only for quantum information processing and simulation [6] but also be helpful in understanding natural processes such as the high efficiency of energy transfer in light harvesting complexes in biological systems [7].

We investigate the behavior of a network of oscillators coupled with external reservoirs as shown in Figure 1. This is a generalization of the Quantum Brownian Motion (QBM) model, which is the paradigm for an open quantum system [1, 8–10]. We present four main original results. First we show that the dynamics is such that: (i) the quantum state always satisfies a simple local master equation and (ii) such equation always has a simple analytical solution. Then we use these results to study the emergence of classical thermodynamical laws in the long time limit. Thus, we derive such laws from first principles in a quantum model by means of a straightforward calculation. Our derivation enables us to show two new thermodynamical results: (iii) We show that it is not possible to create a quantum absorption refrigerator using linear networks. Such refrigerators are interesting systems that, having no movable parts, induce the energy to flow away from a cold reservoir by coupling it with a system that is itself coupled with several other reservoirs. In the quantum regime, absorption refrigerators were recently proposed by Kosloff and others [11, 12] using a specific non-linear model. Our work shows that non-linearity is, in fact, an essential ingredient of a quantum absorption refrigerator. Finally, we show that: (iv) the validity of the third law (unatainability of zero temperature) requires a simple constraint on the type of environmental spectral density, which for low frequencies must behave as ω^α with $\alpha > 0$.



FIG. 1: We analyze the most general network of interacting oscillators coupled with bosonic reservoirs characterized by arbitrary spectral densities and initial thermal states.

We consider a network of N oscillators (in D dimensions) coupled with external environments. Below, we analyze in detail models where the total Hamiltonian is $H_T = H_S + H_E + H_{int}$, where the system Hamiltonian is $H_S = P^T P/2 + X^T V X/2$. The symbol $X = (x_1, \dots, x_K)$ denotes a column vector storing the coordinates ($K = ND$). Similarly, P is a vector collecting all momenta (we consider unit masses and superscript T denotes the transpose). The $K \times K$ matrix V defines the couplings within the system. The environmental Hamiltonian is $H_E = \sum_l H_{E,l}$ with $H_{E,l} = \sum_k ({\pi_k^{(l)}}^2/2m_k + m_k\omega_k^2 {q_k^{(l)}}^2/2)$ and the interaction is $H_{int} = \sum_{l,i,k} C_{ik}^{(l)} x_i q_k^{(l)}$. In what follows, we will study the evolution of the reduced density matrix of the system, obtained by tracing out the environment as $\rho = \mathrm{Tr}_E(\rho_{SE})$.

Before discussing results for models based on the above Hamiltonian it is worth presenting two results that are valid for a larger class of models. In fact, under the single assumption that the evolution preserves the Gaussian nature of the quantum states (together with the hermiticity and trace of ρ) we can show two simple results: (i) $\rho(t)$ always satisfies the following time-local master equation:

$$\begin{aligned}\dot{\rho} \;&=\; -i[H_R(t),\rho] - i\Gamma_{ij}(t)[x_i,\{p_j,\rho\}] \\ &-\; i\tilde{\Gamma}_{ij}(t)[p_i,\{x_j,\rho\}] - D_{ij}(t)[x_i,[x_j,\rho]] \\ &-\; \tilde{D}_{ij}(t)[p_i,[p_j,\rho]] - F_{ij}(t)[x_i,[p_j,\rho]]. \qquad (1)\end{aligned}$$

Here $H_R = P^T M_R^{-1}(t)P/2 + X^T V_R(t)X/2 + f^T(t)X + \tilde{f}^T(t)P$ is a renormalized Hamiltonian that includes time dependent renormalized couplings through $V_R(t)$, a renormalized mass matrix $M_R(t)$ and renormalized forces $f(t)$ and $\tilde{f}(t)$. The non-unitary part of the master equation includes the effect of relaxation (through $\Gamma(t)$ and $\tilde{\Gamma}(t)$) and diffusion (through the matrices $D(t)$, $\tilde{D}(t)$ and $F(t)$). Below, we give formulae for these matrices for the generalized QBM defined by the Hamiltonian described above. Moreover, we can show the following: (ii) The state satisfying equation (1) can be explicitly written as a function of time. For this it is convenient to use the characteristic function, $\chi(\alpha,t)$, which is the expectation value of the displacement: $\chi(\alpha,t) = \mathrm{Tr}(\rho(t)\exp(-i(Pk_p - Xk_x)))$. Here $\alpha = (k_x,k_p)$ is a $2K$–component vector defining momentum and position displacements. We can show that $\chi(\alpha,t)$ is:

$$\chi(\alpha,t) = \chi(\Phi(t)\alpha,0)\exp(-\frac{1}{2}\alpha^T\Sigma_F(t)\alpha)\exp(iJ^T(t)\alpha). \qquad (2)$$

In this equation the time dependence is hidden in the $2K\times 2K-$ matrices $\Phi(t)$ and $\Sigma_F(t)$ and in the $2K$–vector $J(t)$. This results are valid for all linear models (with quadratic Hamiltonians). In such cases (1) and (2) show that the state evolves by a combination of a phase space flow and a Gaussian modulation (as the environment only induces renormalization, friction and diffusion).

Here we sketch the derivation of these results that, to the best of our knowledge, have not been noticed so far. The first result is a generalization of the master equation obtained for QBM in [8]. The second one is a generalization of a well known classical result by Rieder, Lebowitz and Lieb (see [3] for references) who obtained the explicit form of the classical attractor state for general harmonic networks (for QBM the analytic solution was discussed in [10]). Interestingly, both results are valid for arbitrary linear systems and can be obtained analyzing the properties of the evolution super-operator for $\rho(t)$, which is such that $\rho(t) = J(\rho(0))$. Thus, it is simple to show that any super–operator preserving the Gaussian nature of quantum states (together with their hermiticity and trace) can be written in the position representation as: $J(z,z',t;z_0,z_0',0) \equiv \langle z|J(|z_0'\rangle\langle z_0|)|z'\rangle = e^{i(\chi^T b_1 Z + \chi^T b_2 Z_0 + \chi_0^T b_3 Z + \chi_0^T b_4 Z_0)} e^{i(c_1^T\chi + c_2^T\chi_0)} \times e^{-\chi^T a_{11}\chi - \chi^T a_{12}\chi_0 - \chi_0^T a_{22}\chi_0}/2\pi\mathrm{det}(b_3)$. Here we defined K–vectors $Z = (z+z')/2$ and $\chi = z - z'$. The matrices b_m and a_{mn} together with the vectors c_l parametrize any quantum evolution. They depend on the microscopic model, as described below. The way to show the validity of the master equation (1) from the propagator is simple following the method discussed in [1, 9]. We can compute the time derivative of the operator and notice that all the dependence on the initial coordinates z_0 and z_0' can be eliminated by rewriting these terms as a linear combination of others which are proportional to the final coordinates $z^{(\pm)}$ and to the derivatives of the propagator with respect to them. This implies the validity of the master equation (1). Derivation of the solution (2) is even simpler as it can be done by integrating over initial coordinates using the explicit Gaussian propagator.

Coefficients appearing in the equation (1) and in the solution (2) are determined by the time dependent functions appearing in the propagator. They can be obtained from microscopic models such as the one described by the above Hamiltonian. We did this using the Feynman-Vernon method [13] to compute the propagator. It is simple to show that in this case only two properties of the environment determine the propagator: the spectral density and the initial temperatures. The spectral density is $I(\omega) = \sum_l I^{(l)}(\omega)$ where $I_{ij}^{(l)}(\omega) = \sum_k C_{ik}^{(l)}C_{jk}^{(l)}\delta(\omega - \omega_k)/2m_k\omega_k$ (these $K\times K$ matrices are diagonal only if each environment couples with a single site). We assume that the initial state of each environment is thermal with temperature T_l. The spectral density determines the dissipation kernel defined as $\gamma(\tau) = \int_0^\infty d\omega I(\omega)\cos(\omega\tau)/\omega$. The temperature dependent noise kernel is $\nu(\tau) = \int_0^\infty d\omega \sum_l I^{(l)}(\omega)(1+2n_l(\omega))\cos(\omega\tau)$, where $n_l(\omega)$ is the expectation value of the number operator in the l-th environment (i.e., $1+2n_l(\omega) = \coth(\omega/2k_BT_l)$) (we denote $\hat{\nu}(\omega) = \sum_l I^{(l)}(\omega)\coth(\omega/2k_BT_l)$). For this generalized QBM model all coefficients in (1) and (2) depend on the $K\times K$ matrix $G(t)$, which is the solution of:

$$\ddot{G} + V_RG + 2\int_0^t d\tau\gamma(t-\tau)\dot{G}(\tau) = 0. \qquad (3)$$

with initial conditions $G(0) = 0$ and $\dot{G}(0) = I$ (the renormalized potential is defined as $V_R = V - 2\gamma(0)$). The Laplace transform of $G(t)$, denoted as $\hat{G}(s)$, is $\hat{G}(s) = (s^2I + V_R + 2s\hat{\gamma}(s))^{-1}$, where $\hat{\gamma}(s) = \int_0^\infty d\omega I(\omega)s^2/(\omega^2+s^2)$ is the transform of the dissipation kernel. In this model the master equation (1) is such that $M(t) = I$, $f(t) = 0 = \tilde{f}(t)$ (i.e., there is no mass renormalization nor renormalized forces) and $\tilde{\Gamma}(t) = \tilde{D}(t) = 0$ (i.e. there is only normal dissipation and no momentum diffusion). The renormalization and friction matrices are given as $V_R(t) = (\ddot{G}\dot{G}^{-1}\ddot{G} - \dddot{G})(\dot{G} - \ddot{G}\dot{G}^{-1}G)^{-1}$ and $2\Gamma(t) = (\dddot{G}\dot{G}^{-1}G - \ddot{G})(\dot{G} - \ddot{G}\dot{G}^{-1}G)^{-1}$. In turn, the exact solution (2) is such that $J(t) = 0$ and $\Phi(t)$ is

$$\Phi(t) = \begin{bmatrix}\dot{G}(t) & G(t)\\ \ddot{G}(t) & \dot{G}(t)\end{bmatrix}. \qquad (4)$$

For dissipative environments $G(t)$ decays to zero for long times. This is the condition that must be satisfied for the asymptotic state to become independent of the initial state (as seen from (2), this happens when $\Phi(t)$ vanishes). In this asymptotic limit the state is determined by the matrix $\Sigma_F(t)$, which is such that $\Sigma_F(t) = \begin{bmatrix} \sigma_F^{(0,0)}(t) & \sigma_F^{(0,1)}(t) \\ \sigma_F^{(1,0)}(t) & \sigma_F^{(1,1)}(t) \end{bmatrix}$. Each $K \times K$ submatrix can be written as $\sigma_F^{(n,m)}(t) = \int_0^t \int_0^t dt_1 dt_2 G^{(n)}(t_1)\nu(t_1 - t_2)G^{(m)}(t_2)$. Thus, $\Sigma_F(t)$ depends on $G(t)$ and on the temperature-dependent noise kernel. This matrix is related with the diffusive matrices appearing in the master equation. In fact, we can write $D = \mathrm{Sym}(V_R\sigma_F^{(0,1)} + 2\Gamma\sigma_F^{(1,1)} + \dot{\sigma}_F^{(1,1)})$.

The above results enable us to study the long time regime analyzing the emergence of thermodynamical laws. In this case, $\sigma_F^{(n,m)} \rightarrow \mathrm{Re}\int_0^\infty d\omega\omega^{n+m}i^{m-n}\hat{G}(i\omega)\hat{\nu}(\omega)\hat{G}(-i\omega)$. We consider the network to be divided into non–overlapping regions S_l $(l = 1, ..., R)$, each one of which is coupled with an environment E_l (generalization to multiple environments coupled with each region is discussed below). We study the energy transfer between the system and the environments by computing the time derivative of the expectation value of the system's Hamiltonian which can be written as $d(\langle H_R\rangle)/dt = \sum_l \dot{Q}_l$. This equation is equivalent to the first law of thermodynamics, where the heat current entering into the system through the l–th region is $\dot{Q}_l = Tr\left(P_{S_l}V_r\sigma_F^{(0,1)}\right)$, where P_{S_l} is the projector onto the l–th region of the network. In the stationary limit, the conservation law $\sum_l \dot{Q}_l = 0$ is satisfied. Using the formula for $\sigma_F^{(n,m)}$ we find that the heat current is:

$$\dot{Q}_l = \sum_k \int_0^\infty \omega d\omega \dot{\mathcal{Q}}_{lk}(\omega)\coth(\omega/2k_BT_k), \qquad (5)$$

where

$$\begin{aligned} \dot{\mathcal{Q}}_{lk}(\omega) &= \mathrm{Im}\,\mathrm{Tr}(P_{S_l}V_r\hat{G}(i\omega)I^{(k)}(\omega)\hat{G}(-i\omega)), && (6)\\ &= -2\pi|A_l^T(\omega)\hat{G}(i\omega)A_k(\omega)|^2 \qquad j \neq k. && (7) \end{aligned}$$

To derive the last equation we used the explicit form of $\hat{G}(s)$ together with the fact that every spectral density $I^{(l)}(\omega)$ can be written as $I^{(l)}(\omega) = A_l(\omega)A_l^T(\omega)$. The $K-$vector $A_l(\omega)$ has non–vanishing components only for the sites coupled with the environment E_l. In the derivation we also used the condition $\mathrm{Re}(\omega\hat{\gamma}(i\omega)) = \pi I(\omega)$ (which is nothing but the simplest form of the fluctuation dissipation theorem [8]). From these results we find that the heat transfer matrix $\dot{\mathcal{Q}}_{lk}(\omega)$ has the following properties: i) $\dot{\mathcal{Q}}_{lk}(\omega) = \dot{\mathcal{Q}}_{kl}(\omega)$; ii) $\dot{\mathcal{Q}}_{lk}(\omega) \leq 0$ if $j \neq k$; iii) $\sum_l \dot{\mathcal{Q}}_{lk}(\omega) = 0$. Moreover, we also find that the heat flow can be related with the coefficients of the master equation as $\dot{Q}_l \rightarrow \mathrm{Tr}(P_{S_l}(D - 2\Gamma\sigma_F^{(1,1)}))$. Using the above results we can derive the following thermodynamical laws:

a) *Equilibrium: Energy flows through the system if and only if there is a temperature gradient.* This is due to the fact that the heat flow into the j–th site can be written as

$$\dot{Q}_l = -2\sum_{k\neq l}\int_0^\infty \omega d\omega \mathcal{Q}_{lk}(\omega)(n_l(\omega) - n_k(\omega)), \qquad (8)$$

As $\mathcal{Q}_{lk}(\omega) \leq 0$, the heat current $\dot{Q}_l$ vanishes if and only if the $T_l = T_k$ for all l, k. This is equivalent to the 0–th law of thermodynamics: When all the environments have the same temperature then energy does not flow through the system and equilibrium is reached.

b) *Heat flows from the hot to the cold reservoir.* This is nothing but Clausius version of the second law of thermodynamics and can be shown as follows: If T_l is the highest temperature, then $n_l(\omega) > n_k(\omega)$ for all k. Therefore, as $\mathcal{Q}_{lk}(\omega) \leq 0$ we find that $\dot{Q}_l \geq 0$. This implies that the hottest reservoir always injects energy in the system. Of course, the conservation law implies that the other reservoirs absorb this energy. Heat currents are

$$\dot{Q}_l = 4\pi\sum_{k\neq l}\int_0^\infty \omega d\omega|A_l^T(\omega)\hat{G}(i\omega)A_k(\omega)|^2(n_l(\omega)-n_k(\omega)). \qquad (9)$$

It is worth noticing that this same argument can be applied to show that the coldest reservoir always absorbs energy from the system. In fact, if $T_l < T_k$ for all k we have that $\dot{Q}_l \leq 0$. For this reason, we find that it is impossible to build an absorption refrigerator using an harmonic network such as the one we considered. Thus, such type of refrigerator, recently discussed by Kosloff and co-workers [11, 12], requires non-linearity as an essential ingredient. This no-go theorem for the linear absorption refrigerator is general as it applies for any spectral density of the reservoirs.

There are assumptions in the above derivation which can be easily relaxed. Thus, we assumed that each region of the system couples with a single environment. We can generalize this as follows: We assume that the l–th region couples with N_l environments labeled by the index a_l (each having temperature T_{a_l}). For simplicity, we also assume that all the spectral densities $I^{(l,a_l)}$ are the same (this can also be relaxed), i.e. $\sum_{a_l} I^{(l,a_l)}(\omega) = N_l I^{(l)}(\omega)/N_l$. In this case we can show that the heat flow into the region–l is given by equation (9), where the temperature dependent factor $(n_l(\omega) - n_k(\omega))$ must be replaced by a factor $(\tilde{n}_l(\omega) - \tilde{n}_k(\omega))$, where $\tilde{n}_i(\omega) = \sum_{a_i} n_{a_i}(\omega)/N_i$ is the average number of excitations of all the environments coupled with the region S_i. In the high temperature limit, the temperature dependent factor in (9) becomes proportional to the difference between the average temperature of the reservoirs, but this factor depends nonlinearly on the temperatures otherwise. Thus, the second law in this case states that the heat always flows into the region of the system that is coupled with the reservoirs with the largest average number of excitations (or the largest average temperature in the high temperature limit). Equivalently, heat always flows away from the region coupled with the reservoirs with the smallest average number of excitations (temperature). Of course, the environment with the lowest temperature may be coupled with a region which is itself coupled with hotter environments in such a way that the average number of excitations is not the lowest one. Then, heat would flow away from the such environments. However, it is clear that by enlarging the definition of 'the system' and by using the previous argument, one can prove that the environment with the lowest temperature would absorb heat. The other assumption in the above derivation is that different regions S_l do not overlap. However, this can also be generalized using the fact that the obtained results are valid for arbitrary networks. Thus, the case of overlapping regions can be analyzed as follows: Consider a network with regions $S_2 \subset S_1$. If $S_{1,2}$ couple with different environments $E_{1,2}$, this situation is physically equivalent to that of a network where S_1 couples with E_1 while S_2 couples strongly with another network S_2' which is itself coupled with E_2. In such case, all previous results directly apply. As a consequence of this, the no–go theorem for quantum absorption refrigerators applies to all linear networks.

Another version of second law is obtained defining the entropy flow into the system as $\dot{S} = \sum_l \dot{Q}_l/\tilde{T}_l$, where $\tilde{T}_l$ is any monotonically increasing function of T_l. Then $\dot{S} = -\int_0^\infty d\omega \sum_{l\neq k} \mathcal{Q}_{lk}(\omega)(1/\tilde{T}_k - 1/\tilde{T}_l)(n_k(\omega) - n_l(\omega)) \leq 0$.

The above thermodynamical laws are valid for environments with arbitrary spectral densities. Below, we demonstrate that the validity of the third law imposes a constraint on $I(\omega)$. For simplicity, we consider environments coupled with single sites, with spectral densities $I^{(j)}_{jj}(\omega) = \gamma_j \omega^{p_j} \theta(\omega)$ ($\theta(\omega)$ is a cutoff function vanishing when $\omega \geq \Lambda$ and γ_j a coupling constant). When $p_j = 1$ the environment is ohmic while it is super–ohmic (sub–ohmic) for $p > 1$ ($p < 1$). Using this, we can show that: *c) The third law of thermodynamics emerges if* $p_j > 0$.. Thus, the heat flow into the j–th site is:

$$\begin{aligned}\dot{Q}_j &= 4\pi \sum_{k\neq j} \int_0^\infty \omega^{1+p_j+p_k} \gamma_j \gamma_k \theta^2(\omega) d\omega |\hat{G}(i\omega)_{jk}|^2 \\ &\times \sum_{a\geq 1} (e^{-a\omega/k_B T_j} - e^{-a\omega/k_B T_k}). \qquad (10)\end{aligned}$$

For low temperatures the integral is dominated by low frequencies. Taylor expanding the integrand and assuming that all T_j are close to the average $\bar{T}$ we find:

$$\dot{Q}_j = \sum_{k\neq j} \gamma_j \gamma_k \bar{T}^{1+p_j+p_k} (T_j - T_k)(2 + p_j + p_k)\alpha_{kj}. \quad (11)$$

where $\alpha_{lj} = 4\pi(k_B)^{2+p_j+p_k} |\hat{G}(0)_{jk}|^2 \Gamma(2 + p_j + p_k)\zeta(2 + p_j + p_k)$. Thus, for two reservoirs with the same spectral density (i.e., $p_A = p_B$) and a small temperature difference we find that the heat flow is $\dot{Q}_A = 2(p + 1)\gamma^2 \alpha_{AB} \bar{T}^{2p+1} \Delta T$. This implies that the entropy flow into site–A is $\dot{S}_A = \dot{Q}_A/T_A \propto \bar{T}^{2p}$. Therefore, the entropy flow vanishes as a power law when $\bar{T} \to 0$ if $p > 0$, which is nothing but Nernst's version of the third law.

Thus, we presented new results concerning the evolution of open networks of quantum oscillators: (i) we deduced an exact master equation and (ii) we obtained its explicit solution. We used this to show how thermodynamical laws emerge from first principles for general (ohmic and non-ohmic) reservoirs. We proved a no-go theorem for linear absorption refrigerators and showed that the validity of the third law imposes a constraint on the low frequency behavior of the spectral densities. Such constraint is not the same as the one reported in [12]. Thus, according to our results the validity of the third law is consistent with sub-ohmic, ohmic and super-ohmic environments whereas the sub-ohmic ones are excluded according to the [12].

This work was supported by grants from ANPCyT, Conicet and Ubacyt. E.M. is now at Institut für Experimentalphysik, Uni. Innsbruck, Technikerstrasse 25, A-6020 Innsbruck, Austria.

---

[1] J.P. Paz and W.H. Zurek; *Fundamentals of Quantum Information*, Springer, 77-148 (2002).

[2] M. Nielsen and I. Chuang. *Quantum Information and Computation*, Cambridge Univ. Press, Cambridge (2000).

[3] B. Dhar, *Adv. In Phys.* **57**, 457 (2008).

[4] M. Michel, G. Mahler and J. Germer, *Phys. Rev. Lett.* **95**, 180602 (2005); K. Saito, *Europh. Lett.* **61**, 34 (2003). I. Kim and G. Mahler, *Phys. Rev.* **E81**, 1 (2010).

[5] G.D. Lin and L.M. Duan, *New Jour. Phys.* **13**, 075015 (2011).

[6] T. Pruttivarasin, M. Ramm, I. Talukdar, A. Krouter and H. Haffner, *New Jour. Phys.* **13**, 075012 (2011).

[7] M. Mohseni, P. Rebentrost, S. Lloyd and A. Aspriu-Guzik, *Jour. Chem. Phys.* **129**, 174106 (2008).

[8] B.L. Hu, J.P. Paz and Y. Zhang, *Phys. Rev.* **D45**, 2843, (1992).

[9] L. Dávila Romero and J.P. Paz, *Phys. Rev.* **A55**, 4070, (1997).

[10] G. Fleming, B.L. Hu and A. Roura, *Ann. Phys.* **100**, (2011).

[11] A. Levy and R. Kosloff, Phys. Rev. Lett. 108, 070604 (2012)

[12] A. Levy, R. Alicki and R. Kosloff "Quantum absorption refrigerators and the third law of thermodynamics", arXiv:1205.1347 (2012).

[13] R. P. Feynman and F. L. Vernon, *Ann. Phys.* **24**, 118 (1963).